# FISH ASSEMBLAGES OF THE MOROCCAN COASTAL AREA OF AL HOCEIMA (MEDITERRANEAN SEA)

Molinari A., Salvati E., Tunesi L.\* ICRAM, Rome, Italy - a.molinari@icram.org

#### Abstract

The fish assemblage of the marine coastal area of the Al Hoceima National Park has been studied in the framework of a research program for the elaboration of the zoning proposal of the marine park. Collected data show the presence of rich and well diversified assemblages, characterised by a general dominance of small and medium sized specimens.

Keywords: Visual census, Marine Parks, Alboran Sea

#### Introduction

In the framework of the MedMPA project (regional project for the development of Marine and coastal Protected Areas in the Mediterranean region), financially supported by the European commission and coordinated by the Regional Activity Centre for Specially Protected Areas (RAC/SPA-Tunis), ICRAM has been charged with the scientific coordination of the study of the Al Hoceima National Park in Morocco.

Considering the relevance of fish as one of the most important components through which the effects of protection become evident in marine parks [1-3], and the importance of conceiving management measures for fishery and scuba diving activities [4], two surveys were carried out to characterise the coastal fish assemblages in the coastal area applying visual census techniques.

### Material and Methods

The coastline of the Moroccan National Park of Al Hoceima interests about 47 km of Mediterranean coast, characterised by rocky impervious cliffs. The sampling activity was organised subdividing the coastal area into 5 sub-homogeneous coastal units (CU) (Fig. 1), identified on the basis of the main geomorphological features.

Scuba diving paths of 15 minutes were carried out in summer 2002 and 2003 along transects based on four sampling depth ranges (0-3; 4-7; 12-16; 24-30m), in order to acquire semi quantitative data on abundance and size class composition [5]. At least 3 transects were planned perCU. The main sea bottom typologies for each path were registered. A Chi-square test was applied to compare the size class composition recorded in the five considered CUs.




Fig. 1. Study area: the eastward (E) and westward (W) borders of the Al Hoceima National Park are shown. The 5 identified coastal units (I-V) are highlighted.

# Results

The low slope encountered on certain transects sometimes did not allow to reach the 12-16 and the 24-30 m depth ranges. On the whole, 52 paths allowed to identify 69 species (10 Eastern Atlantic and South Mediterranean). Sparids, Labrids and Serranids were the most represented families (15, 14 and 8 species respectively).

Only 17 species occurred in all five CUs while another 17 were each censed only in one CU. This may be due specific characteristics like crypticity (C.conger, P.phycis, S.notata, S.acus), and rarity (L.bergylta, S.cretense, E.aeneus, E.marginatus, P.auriga, S.aurata, T.nobiliana) or because they are associated to soft bottoms (B.podas, Callionymus sp., Gobius sp., S.cinereus, P.erythrinus). The higher number of species were recorded in the costal unit IV (48). Highest species richness were recorded at 0-3 m and 4-7 m (47 species).

Excluding CU I, which is dominated by small specimens (65.8%), and significantly different from the others (chi-square test, p<0.001), the fish assemblage of the marine park was characterised by a general co-dominance of medium sized (from 43.9% to 56.0%) and small specimens (from 33.7% to 43.3%). The large size class is generally

**Table 1. Species recorded in the five coastal units identified along the study area.** (•)) Eastern Atlantic and South Mediterranean species.

Kataly	Research	COLMAND REES	<b>Materials</b>	Nanana.	Chattailled
electricity.	Johanist vis	HILWA	Munterina	Minera creiera	1.17
Apparation	Acades Interfals	KARAWA	Parapertities	- Athereix pinoreix	All
Sto windar	Plantiterritor pallaragine	1.10	Scatidad	Spendown cresionary (	100
	Managements meditarias	300	Son payerstur	Storganu statirnipo	TILW
	Novition to price to 11	an an		disequenci notati:	TV.
	Parahlesolus sous'	8,800,9		Аспервые доком:	1,0
	Farahleenlus compulmiantus:	1.00	Semesidae	Articles perfeite:	URA-
	Scandilla criscita (f.)	III/		Zpinepheliar increas (* )	IV.
Bulledge	Notice chalco	- 19		description make (* )	101
Saltonymilion	- Gallion processor	19		dynapolito nenomino	1
Congrette	- Stragen ranger	10.0		Annaus and edit (* )	1,0
Feggesiden.	Figure all Antonio Marine	100		Saventur zeholin	All
Cadidae	Physic physic	19		Savenue hapation	IUV
Gutelidas	Gelitos Cocchiciali	LWX	CONTRACTOR AND ADDRESS.	Simprove por the	LINE
	Colinos cruchistas	10.03	Sourido.	dioes hopes	All
	Antonic op	78	0.000000	January division	81,016
Dumphou.	Parapetriposes acresimentos-	- 2.1:		Jopinson remains	W/V
Labridau	Contribute	All.		. Opliana nevirus sentine	UUWV
	Chimolebrus superiritis	ILBUVN		-Diplodus purticory	LBUVA
	Labrus iterarilla (*)	0.1E		Stotuday parasty	All
	Latinary remains	E/V		-Hofiston returns	ARI
	Springshould distribute	10		-SMLKE distance	All
	Symphodus dodarists	DURLIN		Jogothar searce	BULWY
	Symphodus mediannesss	All I		Pagellus erysteises	TV.
	Symphodus reglanopricus	BIBLION		/ Reprovisionigal (1)	TV.
	Emphotor recitor	Al Al		Planter water to	URN
	Symphotic acceptant	30		Anger expo	All
	Symphotor microl	Att .		Appenic Andre	IV.
	Symphodus southers	10,000		Spreidylloseme cerebrosi	ULKX
	Symphotoc Street	All	Cymyrel/ddw:	Sympositive execu-	IV
	(Nedepourser service http://	- All	lynoshmiw.	faraute motivies	. 4
Moromico	Megaharather Edward	DIV	front medical	Parallelian discour	19,00
Mingration .	Misprox	30000	ling/terygecon	hydrogowo dodwo	1,10
Militar	Millior Iterature	EW	0.000	Transpolar transminus	All
	Mollar surroumeter	All	110		

less represented (from 8.9% to 15.0%). Taking into account only the first depth range, the I, III and IV CUs were characterised by a high percentage of small specimens (higher than 60%). The higher percentages of large specimens were recorded in the deeper range (24-30 m) in the II and III CUs (23.6% and 24.3% respectively).

## Discussion and Conclusion

The fish assemblage of the Al Hoceima National Park is rich in species and characterised by the relevant presence of Eastern Atlantic and South Mediterranean elements, which testifies the role of the proximate Gibraltar strait. These peculiarities stress the importance of this national park in protecting a very unique Mediterranean fish assemblage in the framework of a network of Mediterranean marine protected areas.

The scarce occurrence of large specimens highlights a relevant fishing pressure, thus emphasizing the importance of implementing new specific protection measures.

Acknowledgments. Prof.S. Benhissoune of the Faculty of Sciences- University of Agadir, Morocco, management of the Ministère des Eauxs et Forêtsand of National Park of Al Hoceima, and the personnel of the I.T.P.M.

# References

1-Villa F., Tunesi L., Agardy T., 2002. Optimal zoning of marine protected areas through spatial multiple criteria analysis: the case of the Asinara Island National Marine Reserve of Italy. Conservation Biology, 16(2): 1-12. 2-Tunesi L., Rais C., Benhissoune S., Agnesi S., Bazairi H., Benhamza A., Di Nora T., Franzosini C., Limam A., Manca Zeichen M., Mo G., Molinari A., Nachite D., Piccione M.E., Sadki I., Salvati E., in press. Zoning of the marine component of the national park of Al Hoceima (Mediterranean, Morocco): a case study. Biol. Mar. Medit., 11. 3-Halpern B.S., 2003. The impact of marine reserve: do reserve work and

3-Halpern B.S., 2003. The impact of marine reserve: do reserve work and does reserve size matter? Ecological Applications, 13 (1) Supplement: S117-S137.

4-Agnesi S., Di Nora T. and Tunesi L.,2001. The study of diving tourism to support the adaptative management in an Italian marine protected area (Ustica island). Rapp. Comm. Int. Mer. Médit., 36: 347.
5-Harmelin-Vivien M.L., Harmelin J.G., Chauvet C., Duval C., Galzin R., Lejeune P., Barnabe G., Blanc F. and Chevalier R., 1985. The underwater observation of fish communities and fish populations: Methods and problems. Rev. Ecol. Terre Vie, 40: 467-540.

Rapp. Comm. int. Mer Médit., 37,2004